

# **HOW TO USE**

**DUAKUR® PRESSAIR®**INHALER FOR COPD

## 1. **READY** - BEFORE INHALATION



Remove the protective cap by lightly squeezing the arrows marked on each side and pulling outwards.



Hold the **DUAKLIR®**PRESSAIR® inhaler with
the mouthpiece toward
you and the orange
button facing straight up.
Do not tilt.



Press the orange button all the way down and release to prepare your dose. Do not tilt.



Do not hold the orange button down. Stop and check to make sure the dose is ready for inhalation.



Check to see if the color of the control window has changed from red to green. If the controller window remains red, go back to Step 1.



The green control window confirms that your medication is ready for inhalation.

## 2. CLICK - DURING INHALATION



Hold the inhaler away from your mouth, and breathe out completely. Never breathe into the inhaler.



Put your lips tightly around the mouthpiece of the **DUAKLIR®** PRESSAIR® inhaler. Take a strong, deep breath. A "click" sound will occur during inhalation when the inhaler is used correctly. Keep breathing in as long as possible, even after you have heard the "click",\* to be sure you get the full dose. Remove the **DUAKLIR®** PRESSAIR® inhaler from your mouth and hold your breath as long as possible. Then slowly breathe out, away from the inhaler.

### 3. CONFIRM - AFTER INHALATION



Make sure the control window has turned green to red. This "confirms" that you've inhaled your full dose correctly. If the window did not change to red, you may have forgotten to remove your finger from the orange button before inhaling, or you may not have inhaled correctly. If this happens, go back to Step 2 and repeat the dosing instructions.



Once the control window turns red, place the protective cap back onto the mouthpiece of the **DUAKLIR®** PRESSAIR® inhaler.

\*Note: Some people may not hear the "click." Use the control window to ensure you have inhaled correctly

### INDICATION AND USAGE

**DUAKLIR®** PRESSAIR® is a prescription medicine to treat chronic obstructive pulmonary disease (COPD). COPD is a long-term (chronic) lung disease that includes chronic bronchitis, emphysema or both.

**DUAKUR®** PRESSAIR® is not for the treatment of asthma.

DUAKUR® PRESSAIR® is not a rescue medicine and should not be used for treating sudden breathing problems.

Always have a rescue inhaler with you to treat sudden breathing problems. Get emergency medical care if your breathing problems worsen quickly or you use your rescue inhaler, but it does not relieve your breathing problems.

Please see additional Safety Information below and the accompanying Full Prescribing Information at duaklir.com

## **USEFUL TIPS** - Help Manage COPD Over the Long Term



## STICK TO YOUR TREATMENT PLAN

Follow your doctor's advice on how and when to take medication. Remember to always use the inhaler correctly for best results.



## GET REGULAR CHECKUPS

Schedule a few visits with your doctor every year - even if you feel fine! Checking in can help you know if you're on the right treatment plan.



Make sure you stay active and eat right. It can be good for your body. A simple way to start is to take walks and eat a balanced diet.



If you find it difficult, reach out to a doctor for a professional recommendation.

#### IMPORTANT SAFETY INFORMATION

### WHO SHOULD NOT USE DUAKLIR® PRESSAIR®?

Do not use DUAKLIR® PRESSAIR® if you have asthma without also using a medicine called an inhaled corticosteroid.

Do not use DUAKLIR® PRESSAIR® if you have a severe allergy to milk proteins, or are allergic to aclidinium bromide, formoterol fumarate or any of the ingredients in DUAKLIR® PRESSAIR®. Ask your healthcare provider if you are not sure.

#### WHAT SHOULD I TELL MY HEALTHCARE PROVIDER BEFORE USING DUAKUR® PRESSAIR®?

Tell your healthcare provider about all your medical conditions, including heart problems, high blood pressure, seizures, thyroid problems, diabetes, eye problems (especially glaucoma), prostate or bladder problems or problems passing urine, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed.

Tell your healthcare provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, herbal supplements and eyedrops. **DUAKUR®** PRESSAIR® and certain other medicines may interact with each other and may cause serious side effects. Especially tell your healthcare provider if you take anticholinergics, other LABA (long-acting beta²-adrenergic agonist) medicines, xanthine derivatives, steroids, diuretics, monoamine oxidase inhibitors, tricyclic antidepressants, beta-blockers or atropine.

Use **DUAKUR®** PRESSAIR® exactly as prescribed. Do not use **DUAKUR®** PRESSAIR® more often than prescribed or take more medicine than prescribed for you.

### WHAT ARE THE POSSIBLE SIDE EFFECTS OF DUAKUR® PRESSAIR®?

Using too much of a LABA medicine may cause chest pain, increased blood pressure, fast and irregular heartbeat, headache, tremor, or nervousness.

Sudden breathing problems may occur immediately after use of **DUAKLIR®** PRESSAIR® and may be life threatening. If you have sudden breathing problems immediately after inhaling **DUAKLIR®** PRESSAIR®, stop using **DUAKLIR®** PRESSAIR® and call your healthcare provider right away or go to the nearest hospital emergency room.

Call you healthcare provider or get emergency medical care if you have any symptoms of a serious allergic reaction including rash, swelling of the face and tongue, hives, or breathing problems.

You may experience high blood sugar (hyperglycemia) or low blood potassium (hypokalemia). Low blood potassium may cause muscle spasms, muscle weakness, or abnormal heart rhythm.

These are not all the possible side effects of **DUAKUR®** PRESSAIR®.

### ADVERSE REACTIONS

You may report an adverse event related to Covis Pharma products by calling 1-877-411-2510 or emailing <u>medinfoUS@covispharma.com</u>. If you prefer, you may contact the U.S. Food and Drug Administration (FDA) directly at fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Safety Information above and the accompanying Full Prescribing Information at duaklir.com

